

Attachment to Serial No. 09/253,573



Subscribe today
Subscribe for a FREE issue

NewScientist.com

HOME | NEWS | HOT TOPICS | THE LAST WORD | OPINION | WEELINKS | PRINT EDITION | SUBSCRIBE | ARCHIVE | JOBS & CARE



ALL LATEST NEWS

SEARCH NEWS



TOP NEWS STORIES

Facial tumours kill off
Tasmanian devils

Cellphone software
allows a musical jam

Satellite data reveals
rapid Arctic warming


Stem cell mobility
linked to cancer's
spread

Clay's matchmaking
could have sparked
life

Laser system to
pinpoint snipers fast-
tracked

All latest news

FOR SUBSCRIBERS



- Subscribe
- Renew
- Change address

TOP SCIENCE JOBS

Best live jobs in the
hottest areas of
sciences

Materials Science

Bioinformatics

The World's No.1 Science & Technology News Service

'Miracle' gene therapy trial halted

14:30 03 October 02

NewScientist.com news service

A "miracle" gene therapy treatment for children suffering from the fatal "bubble boy" disease has been halted in France, after one of the patients developed leukaemia as a direct consequence of the treatment. However, British doctors argue that without the treatment many of the patients are certain to die, and say a similar trial in UK will continue.

Boys with X-SCID (Severe Combined Immunodeficiency) have a faulty copy of a gene on their X chromosome that makes an immune protein called interleukin-2. As a result, they have no resistance to infection and die unless treated.

In 2000, a team led by Alain Fischer at Necker Hospital, Paris, carried out the first gene therapy treatment, which replaced the faulty gene. It was one of only a handful of successful gene therapy trials in people. In April 2002, the mother of a Welsh boy treated at Great Ormond Street hospital in London described his progress as "nothing short of a miracle".

A total of 15 patients, have been treated so far - 11 in Paris and four in London. But it was revealed on Thursday that one three-year-old patient has now developed leukaemia.

Uncontrolled production

The boy underwent gene therapy at the age of six months, and contracted chicken pox at two-and-a-half. His white cell count increased in response to the infection, as would be expected. But his bone marrow then started uncontrollably producing these cells.

Gene therapy involves shuttling the gene into a patient's cells using a harmless virus. But transferred genes cannot be targeted to insert into a specific part of a chromosome. And it appears, scientists say, that in this boy the new gene was inserted next to an oncogene, called Lmo2, triggering the leukaemia.

This kind of severe side-effect was not unanticipated. "In its original review of these studies, the UK Gene Therapy Advisory Committee recognised the possibility of cancer occurring as a result of

Subscribe to New Scientist for more news and features

Related Stories

Gene therapy cures "bubble boy"



SCIENCE NEWS

All the best science & web

Scientists discover 5 fish in billion-dollar tr world's oceans

The Independent, UK

Google eyes \$15bn c Financial Times, UK

Worms hold 'eternal' BBC

Scientists find first di tumor Reuters

Clinton deal to get Ai world's poor Guardian Unlimited,

Museum claims fossil vertebrate Washington Post

Last fare-paying Con lands in New York CNN

09/253, 373

Pharmacology
Synthetic Chemistry
Regulatory Affairs
Or search all of our
jobs and opportunities

Sponsored Links

Heat Kills Cancer
Aggressive
treatment avail in
U.S. Proven results
for many cancers
www.ichtpt.com

**New CLL
Treatment
Options**
Learn about the
latest advances for
CLL patients
dealing with a
relapse
www.healthtalk.com

**Do you have
colon cancer?**
Investigational drug
clinical trial Need
patients with
advanced disease
<http://etrialis.novartis.com/>

**Cancer
Diagnosis?**
10 secrets to
survival you must
know within one
week of diagnosis.
www.survivors-wisdom.com

this type of gene therapy,"
says Norman Nevin, GTAC's
chairman.

But Nevin confirmed that the
UK trials will continue: "GTAC
is satisfied that all parents and
children treated were
informed of this risk and
received appropriate
counselling prior to
treatment." The chances of
the gene being inserted next
to an oncogene are very low,
he adds.

Some children with X-SCID
are candidates for bone
marrow transplant treatments, but others will undoubtedly die
without gene therapy. Doctors at Great Ormond Street say two
UK patients have died in 2002, because they did not start gene
therapy in time. "It is ethically justifiable to go ahead with the
Great Ormond Street trials because of the benefits accrued from
the treatment," says Nevin.

The full investigation into the three-year-old's leukaemia will take
between 12 and 18 months. Doctors say he is responding well to
chemotherapy.

Emma Young

3 April 2002

Gene therapy causes cancer in mice
18 April 2002

Master switches to boost gene
therapy

3 September 2001

For more related stories
search the print edition Archive


Weblinks

Hôpital Necker-Enfant Malades

SCID, Great Ormond Street Hospital

Gene Therapy Advisory Committee

NewScientist



ADVERTISEMENT

For unbiased reviews
cameras use DigitalC

Digital Camera
Digital Cameras
Digital Camera Review
Digital Camera Review
Sony Digital Camera
Canon Digital Camera
Camcorder
Camcorders
Canon 300D
Digital Camera Links

Print this article Send to a friend

SUBSCRIBE AND SAVE 79%

Get 51 issues for only \$51
saving 79% off the single issue price



First Name
Last Name
Address

City
State
Zip
USA
Email

If you are not from the US click here

Send me New Scientist at 79% off